

ENDOWING SINGLE POLYMER CHAINS WITH ENZYME-MIMETIC ACTIVITY
AND SELECTIVITYJOSÉ A. POMPOSO^{1,2,3}¹*Centro de Física de Materiales (CSIC, UPV/EHU)-Materials Physics Center, Paseo Manuel de Lardizabal 5, E-20018 San Sebastián, Spain*²*Departamento de Física de Materiales, Universidad del País Vasco (UPV/EHU), Apartado 1072, E-20080 San Sebastián, Spain*³*IKERBASQUE - Basque Foundation for Science, Alameda Urquijo 36, E-48011 Bilbao, Spain – Email: Josetxo.pomposo@ehu.es*

Abstract

Linear polymer chains can be folded / collapsed to individual, single-chain nanoparticles (SCNPs) by means of reversible^{1,2} and irreversible³ intrachain crosslinking techniques. SCNP formation is reminiscent of protein folding (see Figure 1), although current synthetic methods lack the perfection of protein folding to functional, globular enzymes.⁴ Nevertheless, in recent years the classical structure–function paradigm (*i.e.*, amino acid sequence → 3D structure → function) has been revisited by taking into account that many non-structured segments of proteins, and even totally disordered proteins, play important roles in protein function.⁵ In this lecture we highlight the significant added value (enzyme-like catalysis, drug binding and delivery) that can be endowed to SCNPs by taking inspiration from the functions of both ordered (native) and disordered proteins.⁶ Specifically, the recently reported “concurrent” binding/folding strategy⁷ opens new, promising avenues for endowing SCNPs with enzyme-mimetic activity and selectivity.⁸




Figure 1. Illustration of hard (a) and soft (b) synthetic nano-objects. Biomimetic SCNPs can be constructed by taking inspiration from the functions of soft, natural nano-objects (c) and, in particular, from both native and intrinsically disordered proteins.⁶

¹ Sanchez-Sanchez, A.; Fulton, D. A.; Pomposo, J. A. *Chem. Commun.* **2014**, 50, 1871-1874.

² Sanchez-Sanchez, A.; Pomposo, J. A. *Part. Part. Syst. Charact.* **2014**, 31, 11-23.

³ Sanchez-Sanchez, A.; Perez-Baena, I.; Pomposo, J. A. *Molecules* **2013**, 18, 3339-3355.

⁴ Altintas, O.; Barner-Kowollik, C. *Macromol. Rapid Commun.* **2012**, 33, 958-971.

⁵ Tompa, P.; Han, K. H. *Phys. Today* **2012**, 65, 64-65.

⁶ Pomposo, J. A. *Polym. Int.* **2014**, 63, DOI: 10.1002/pi.4671.

⁷ Perez-Baena, I.; Barroso-Bujans, F.; Gasser, U.; Arbe, A.; Moreno, A. J.; Colmenero, J.; Pomposo, J. A. *ACS Macro Lett.* **2013**, 2, 775-779.

⁸ Sanchez-Sanchez, A.; Arbe, A.; Colmenero, J.; Pomposo, J. A. *ACS Macro Lett.* **2014**, 3, Submitted.